

Role of Urban Forestry in Improving Air Quality and Human Health

Israr Ullah*¹, Muhammad Ahmad², Muhammad Abbas Khan³, Samiullah Memon⁴,
Phulan⁵, Ikram Ullah⁶, Muhammad Akram⁷, Tauseef Iqbal⁸

¹ College of Agriculture, University of Sargodha, Punjab Pakistan.

Corresponding author: Ullahisrar265@gmail.com

² Department of Forestry and Range Management, Bahauddin Zakariya University, Multan. 66000, Pakistan. mahmadofficialfrw@gmail.com

³ Department of Horticulture, Balochistan Agriculture College, Quetta. mhammadabbaskhan1121@gmail.com

⁴ School of Architecture, Tianjin University, Tianjin, China. ar_samiullah45@yahoo.com

⁵ Department of Environmental science, Government College University Faisalabad phulantapi@gmail.com

⁶ Forestry and Rang Management Faculty of Agriculture, University of Agriculture Faisalabad ikramullah1130@gmail.com

⁷ Department of forestry and range management, university of Agriculture Faisalabad 2020ag2890@uaf.edu.pk

⁸ School of Agricultural Sciences, Zhengzhou University, Zhengzhou, China tauseefiqbal@gs.zzu.edu.cn

DOI: <https://doi.org/10.63163/jpehss.v3i4.920>

Abstract

Urban forests serve as critical green infrastructure for mitigating air pollution, enhancing human health, and promoting climate resilience in rapidly urbanizing environments. This review synthesizes biophysical mechanisms of pollutant removal, including direct absorption of gaseous pollutants and interception of particulate matter (PM), alongside indirect benefits such as Urban Heat Island (UHI) mitigation through shading and evapotranspiration. It addresses challenges like biogenic volatile organic compound (BVOC) emissions, which can exacerbate ozone formation, emphasizing strategic species selection and atmospheric modeling for optimization. Quantitative valuation using tools like i-Tree and the Value of a Statistical Life (VSL) framework demonstrates substantial economic returns, including billions in health savings from reduced morbidity and mortality. Epidemiological evidence links urban greenery to lower mortality risks, improved mental health, and immunological boosts via phytoncides. The paper advocates for equitable policy integration, overcoming barriers like infrastructure conflicts and pollution stress, to ensure resilient, multifunctional urban forests as essential public health assets.

Keywords: Urban Forestry, Air Quality, Human Health, Green Infrastructure, Pollutant Removal, Urban Heat Island (UHI), Biogenic Volatile Organic Compounds (BVOCS), Ecosystem Services, Climate Resilience, Environmental Equity

I. Introduction


The pervasive trend of global urbanization, characterized by sustained population growth and economic development, has conferred numerous societal benefits, including advanced healthcare, expanded employment opportunities, and improved communication networks. However, this rapid

expansion, particularly in developing nations, has concurrently driven significant negative externalities (Nowak et al., 2014). (Humbal et al., 2023). Unplanned urbanization results in land cover changes, the expansion of impervious surfaces, natural resource depletion, and substantial environmental degradation, imposing significant costs on global society (Thompson et al., 2024). The consequence for residents is concentrated exposure to two primary public health threats: pervasive air pollution, including particulate matter (PM), ozone (O₃), and nitrogen oxides (NO_x), and the Urban Heat Island (UHI) effect, which severely modifies local energy balance (Escobedo et al., 2011). Addressing these challenges requires moving beyond traditional point-source pollution control methods toward mitigating human exposure at the receptor level. This imperative places the urban forest the entirety of trees and vegetation within a city at the center of public health strategy, necessitating a rigorous review of its role as a nature-based solution for atmospheric remediation and health promotion. The collective urban forest of the United States alone is quantified to remove approximately 711,000 tonnes of criteria air pollutants per year, providing a direct basis for the valuation of avoided mortality and morbidity benefits (Markevych et al., 2017).

Urban forestry is defined not merely as passive green space maintenance, but as the active care and strategic management of single trees and tree populations within urban settings specifically for the purpose of improving the urban environment. It is an integrated, interdisciplinary practice that involves both meticulous planning and systematic management, including the programming of long-term care and maintenance operations. A defining characteristic of urban forestry is its intrinsic multifunctionality (Tong et al., 2020). It yields a vast array of ecosystem services simultaneously, spanning economic, environmental, and sociocultural benefits. These services range from providing crucial habitats and shelter for wildlife to delivering essential connectivity for biodiversity and significant human health benefits. Critically, successful urban forestry advocates for the recognition of the urban tree canopy as essential urban infrastructure. This reclassification is crucial for strategic policy implementation. Trees, through mechanisms like dry deposition, remove pollutants from the atmosphere, providing direct health co-benefits that justify their strategic implementation within densely populated urban environments (Abhijith et al., 2017).

When trees are viewed purely as an aesthetic amenity, funding is often discretionary and inconsistent. By contrast, classifying the urban forest as critical infrastructure analogous to road networks or utility systems allows municipal governments to justify strategic, long-term capital investment for management, maintenance, and expansion. This elevates urban forestry to a non-negotiable component of essential city operations, providing sustained justification for comprehensive management plans that ensure the urban forest is resilient and can deliver its full range of ecosystem services effectively (Selmi et al., 2016). Furthermore, recognizing the green space-health nexus requires exploring multiple pathways, including air quality mediation and the psychological restoration capacity enabled by reduced environmental stressors (Idris et al., 2025).

Figure 1.1 Role of Urban forestry in Improving Air Quality and Human Health



2. Biophysical and Chemical Mechanisms of Atmospheric Remediation

Urban forests modulate atmospheric quality through sophisticated biophysical and chemical interactions, which can be broadly categorized into direct removal mechanisms and indirect atmospheric modifications. Table 1.1 summarizes the two main ways urban forests directly and indirectly modify the atmosphere, including the pollutants affected and the long-term management required.

Table 1.1 Biophysical Mechanisms of Atmospheric Remediation

Mechanism Category	Specific Mechanism	Pollutants/Effects Addressed	Process/Details	Management Consideration
Direct Removal	Gaseous Absorption	Ozone (O ³), Nitrogen Oxides (NO _x), Sulfur Dioxide (SO ₂)	Pollutants are absorbed through stomata during photosynthesis	High pollutant levels can compromise tree health and reduce filtering capacity
Direct Removal	Particulate Interception	Particulate Matter (PM), especially PM ^{2.5}	Particles "catch" and deposit on leaf/stem surfaces Efficiency is higher with large Leaf Area Index (LAI) and complex leaf microstructures	PM retention is temporary ; requires integrated ground management (permeable pavements) to ensure permanent sequestration (prevent resuspension).
Indirect Modification	Urban Heat Island (UHI) Mitigation	High City Temperatures Energy Consumption	Achieved through shading and evapotranspiration lowering air conditioning needs by up to 30%	Reduces fossil fuel burning for electricity, leading to avoided emissions of pollutants and greenhouse gases

Direct Pollutant Removal: Capture and Absorption Kinetics

Gaseous air pollutants, including ozone (O₃), nitrogen oxides (NO_x), and sulfur dioxide (SO₂), are primarily removed from the atmosphere via absorption through the stomata of the tree foliage during normal photosynthetic gas exchange (Thompson et al., 2024). This biological process is highly effective under optimal conditions, leading to a significant net reduction in urban ozone formation, a finding supported by multiple studies conducted by the USDA Forest Service (Viippola et al., 2018). However, the efficacy of gaseous removal is highly sensitive to the concentration and chemical identity of the pollutants themselves.

High ambient levels of pollutants, particularly O₃ and NO₂, can induce specific phytotoxic responses in plants. These injuries include foliar damage, the loss of pigments, premature senescence (aging), and a decrease in photosynthetic and growth rates. This creates a negative feedback loop: the greater the pollution, the more the trees' health is compromised, consequently reducing their capacity to filter air, thereby diminishing the long-term effectiveness of the urban forest as a pollution sink (Roman et al., 2018). This phenomenon underscores the need for localized assessments of pollutant stress to ensure the long-term viability of the urban forest as an environmental remediator.

Urban forests are highly effective at intercepting airborne particulate matter (PM), including the highly hazardous fine particulate matter (PM_{2.5}). Trees remove these particles by acting as temporary filters, "catching" them as they are deposited onto the surfaces of leaves and stems. This physical interception process is strongly influenced by the structure of the canopy, a critical design parameter for urban planning (Nowak et al., 2013). The efficiency of PM removal is directly correlated with the morphological and physiological features of the vegetation. Specifically, trees with an extended total leaf area (high Leaf Area Index, or LAI) and complex microstructures on the leaf surface tend to promote greater PM deposition, maximizing interception kinetics.

Once deposited, the particulates remain clinging to the surfaces. Crucially, PM retention by trees is often temporary. Particulates can be resuspended into the atmosphere, particularly by wind, allowing them to re-enter the breathable air. Alternatively, precipitation events (rain or snow) dissolve the particulates, transferring them into the stormwater runoff or directly into the soil (Chen et al., 2021). Quantitative modeling reinforces this understanding, with individual urban trees in certain European case studies estimated to remove 0.7–1.4 kg of PM per year, a benefit which requires permanent ground sequestration to prevent resuspension. (Idris et al., 2025).

For the air quality benefit to be maximized and sustained, urban forestry must be managed in tandem with integrated land management. This requires combining specific tree species optimized for PM interception with comprehensive ground-level strategies, such as permeable pavements and soil management, to ensure the permanent sequestration of washed-off PM, preventing its return to the atmosphere (Baró et al., 2014). Empirical measurements utilizing advanced techniques like eddy covariance in urban micro-environments have confirmed the localized efficacy, demonstrating PM reduction rates between 20–30% within urban street canyons. The temporary nature of PM deposition creates a crucial causal link, mandating co-management between green infrastructure (trees) and ground-level grey infrastructure (permeable pavements/stormwater management) to ensure the permanence of the pollution removal benefit.

Indirect Atmospheric Modification and Climate Regulation

The most significant indirect air quality benefit provided by urban forestry is its role in mitigating the Urban Heat Island (UHI) effect. The conversion of natural pervious surfaces to heat-absorbing impervious urban structures significantly alters the local energy balance, driving up city temperatures (Tzoulas et al., 2007). Urban forests combat the UHI effect through two primary mechanisms: shading and evapotranspiration. This provides essential climate regulation services that are becoming increasingly vital in the face of progressive global climate change and the rising frequency of extreme heat days. The i-Tree software tools developed by the USDA Forest Service allow for comprehensive national-scale health and climate valuations of these benefits, translating ecological function into policy-relevant metrics. (Thompson et al., 2024)

Furthermore, shading provided by the urban canopy directly reduces the need for mechanical cooling in buildings, lowering air conditioning needs by up to 30 percent. This reduction in energy consumption correspondingly decreases the amount of fossil fuels burned to produce electricity (Kabisch et al., 2016). This, in turn, achieves significant avoided emissions of criteria air pollutants and greenhouse gases. The cooling benefit, which is external to the tree's biological health and stomatal conductance constraints, offers a reliable, indirect pollution control mechanism. This combination of direct carbon storage in wood, carbon removal from the atmosphere, and the cooling effect makes urban trees exceptionally efficient tools for pollution and climate mitigation (Roy et al., 2012). For instance, the carbon storage capacity of urban and community trees in the United States alone is estimated at 706 megatonnes (Mt).

The UHI mitigation provides a dual benefit: direct thermal stress reduction for human health and indirect pollution control through energy savings. The magnitude of this indirect benefit reinforces

the necessity of viewing urban forestry as bundled infrastructure that solves climate (C storage), energy (cooling savings), and health (VSL) problems simultaneously.

3. Navigating the BVOC Trade-Off: Optimized Species Selection for Air Quality

The relationship between urban forests and air quality is complexified by the biogenic volatile organic compound (BVOC) emissions profile of certain tree species. Addressing this nuance requires moving beyond generalized greening efforts to highly strategic, chemistry-informed planning (Eisenman et al., 2019).

The Atmospheric Chemistry of Biogenic Volatile Organic Compounds (BVOCs)

Trees naturally emit various BVOCs, primarily isoprene and terpenes, into the atmosphere. These compounds are crucial factors in biosphere-atmosphere interactions and atmospheric chemistry. However, BVOCs are highly reactive and act as precursors to the formation of secondary air pollutants. When BVOCs are emitted in the gaseous phase, they are oxidized in the atmosphere, forming more functionalized compounds that can condense into secondary organic aerosols, contributing to fine particulate matter (PM_{2.5}). More significantly, in urban areas characterized by heavy traffic and industrial activity, the environment is typically rich in nitrogen oxides (NO_x), placing it in a "VOC-limited" photochemical regime (Twohig-Bennett & Jones, 2018).

Under these high NO_x conditions, the oxidation of BVOCs reacts strongly with NO_x to significantly enhance the formation of ground-level ozone (O₃). This contribution of BVOCs to ozone formation challenges the simple assertion that all urban greening uniformly improves air quality, particularly because O₃ is a severe respiratory hazard (Thompson et al., 2024). Moreover, the emission rate of BVOCs is not static; it is influenced by environmental conditions, with higher temperatures and existing pollution stress potentially exacerbating BVOC emissions (Gascon et al., 2016). This interdependence necessitates advanced atmospheric modeling to predict the net local air quality impact, demanding precise species selection. (Idris et al., 2025).

Strategic Management through Species Optimization

The potential for BVOC emissions to adversely impact O₃ and PM_{2.5} air quality necessitates that urban forestry planning shift its focus from maximizing total tree canopy to optimizing species composition. This requires interdisciplinary collaboration between urban foresters, atmospheric scientists, and public health officials, making species selection an exercise in localized atmospheric modeling. Policy mandates for urban greening efforts must now include considering the BVOC emission potential of candidate species, treating species selection as a critical risk management function (Konijnendijk et al., 2023).

Scientific assessments have resulted in qualitative ratings—low, medium, or high—assigned to species based on their ozone and PM_{2.5} formation potential. The difference in impact between low and high emitters is substantial, illustrating the importance of precision in urban ecology planning. Researchers have developed standardized tools, such as the urban tree air quality score, to rank trees based on their potential net effect on urban air quality (Dadvand et al., 2022). Therefore, policy recommendations must strongly advocate for a shift from planting based purely on horticultural ease or aesthetics to planting based on atmospheric chemistry modeling, ensuring that the infrastructure investment consistently yields a net positive outcome for human health.

The BVOC trade-off elevates urban forestry from a general 'greening' initiative to a field requiring highly specific, model-driven, and local-scale atmospheric engineering. This technical requirement mandates the use of atmospheric transport models (like those used in i-Tree or specialized local models) before deployment to ensure the net air quality benefit is positive, mitigating the risk of photochemical smog generation. (Idris et al., 2025).

4. Quantification and Valuation: Establishing the Return on Investment (ROI)

To secure political feasibility for large-scale urban forestry initiatives, abstract ecological and health benefits must be translated into quantifiable, monetized financial metrics, establishing a verifiable Return on Investment (ROI). This translation is necessary to allow green infrastructure to compete effectively against traditional grey infrastructure expenditures (McPherson et al., 2022).

Modeling Ecosystem Services and Pollutant Removal

The i-Tree software suite, developed by the USDA Forest Service, provides a standardized, peer-reviewed framework for assessing and valuing the urban forest resource. Tools like i-Tree Eco and i-Tree Design facilitate data collection (via inventory or field evaluation) to quantify the dollar value of annual environmental benefits. This modeling framework is essential for overcoming implementation barriers that often plague urban nature-based solutions (Nowak et al., 2014).

i-Tree models quantify several critical benefits: air pollution removal (tonnage of hazardous pollutants intercepted), carbon sequestration and storage, and energy conservation (savings from reduced energy demand due to shading). The quantification of these benefits is vital for providing political and financial justification. Global assessments, which systematically review urban tree benefits, costs, and assessment methods across different climatic zones, confirm that proactive, strategic planning is universally justified, demonstrating strong cost-benefit ratios across cities (Escobedo et al., 2011).

Monetizing Health Outcomes: The VSL Framework

The most impactful economic metric for validating environmental policy is the quantification of health benefits derived from reduced mortality risk. This is achieved using the Value of a Statistical Life (VSL) framework. The VSL reflects the aggregated willingness-to-pay by a large population for small reductions in their individual risk of death from environmental causes (Markevych et al., 2017). The application of VSL provides policymakers with the methodology to quantify the economic benefits of avoided fatalities caused by improved air quality. The calculation of the ROI is crucial for political viability.

The monetary evidence is robust: meta-analyses summarizing the financial impact of air quality improvement show that urban forests generate an estimated 6.8 billion per year in US health savings derived from reduced morbidity and mortality. This significant economic benefit is directly linked to the removal of hundreds of thousands of tonnes of criteria pollutants annually, as established by national-scale modeling (Tong et al., 2020). The economic justification for urban forestry must be viewed as bundled infrastructure that solves climate (C storage), energy (cooling savings), and health (VSL) problems simultaneously. (Idris et al., 2025).

5. Multi-Pathway Human Health Impacts and Immunological Benefits

The benefits of urban forestry extend beyond atmospheric chemistry, providing a synergistic defense against chronic and acute health risks. The overall net benefit of greenspaces is derived from a complex range of environmental, social, psychological, and physiological effects.

Epidemiological Evidence: Reduced Morbidity and Mortality

Robust epidemiological evidence suggests that diminished exposure to air pollution is a primary reason for the salutary effects of urban greenness on human health. By reducing ambient concentrations of pollutants, urban forests mitigate the risk of adverse cardiovascular and respiratory outcomes (Abhijith et al., 2017). Systematic reviews confirm that increased residential greenspace exposure is associated with a reduction in mortality risk.

Specific analyses, aggregating 20 studies, demonstrate a consistent finding of an approximate 12% reduction in overall mortality associated with higher levels of nearby green space. These health gains are compounded by increased proximity to green space, which facilitates healthier behaviors, such as increased physical activity (Thompson et al., 2024). Furthermore, trees contribute significantly to acoustic regulation, reducing unwanted sounds from traffic and other sources, which is important given the negative impact of environmental noise on cardiovascular health (Selmi et al., 2016). People often perceive noise reduction even from sparse planting strips, indicating that the visual and acoustic aspects of urban vegetation interact to positively alter the perception and evaluation of soundscapes.

Finally, the mitigation of the UHI effect through shading and evapotranspiration serves as a critical, acute health defense mechanism. By preventing excessive temperature exposure, urban forests directly reduce heat-related morbidity and mortality, an increasingly important function given global climate change (Viippola et al., 2018). Cohort studies link greenspace proximity to quantifiable longevity benefits, showing a gain of 0.5 years in life expectancy, offering a profound validation of the public health investment.

Restoring Capacity and Active Health Promotion (The Phytoncide Pathway)

Urban forests provide crucial mechanisms for restoring mental capacity and promoting active physiological health. Studies have demonstrated the positive impact of urban green space on mental well-being, including reduction of perceived stress, improved sleep quality, and general mental health restoration. These psychological benefits comprise a significant portion of the total health impacts associated with tree exposure (Roman et al., 2018). A profound and often underappreciated health mechanism involves chemical ecology: the emission of biogenic amines, specifically phytoncides.

These organic compounds, which possess natural antibacterial and antifungal qualities that help plants fight disease, are inhaled by humans exposed to the forest environment. When these chemicals enter the human body, the immune system responds by increasing the number and activity of Natural Killer (NK) cells. NK cells are a type of white blood cell that specifically targets and eliminates tumor- and virus-infected cells (Nowak et al., 2013). The sustained nature of this immunological boost is significant; research has indicated that increased NK cell activity resulting from short forest bathing trips can persist for more than 30 days. This suggests that urban forest exposure can be strategically framed as a preventative immunological treatment. The literature confirms this broad range of mechanisms and recommends integration strategies based on these holistic benefits.

The significant epidemiological evidence for mortality reduction (12%) and gains in life expectancy (0.5 years) suggests a profound synergistic effect where environmental quality mediators (clean air, cooling) combine with innate biological stimulants (phytoncides) and psychological restoration. This combined effect establishes urban forestry as a comprehensive, layered public health intervention that simultaneously remediates the environment and proactively enhances human biological resilience (Chen et al., 2021).

6. Strategic Implementation, Policy Integration, and Environmental Equity

The strategic deployment and long-term maintenance of urban forests face substantial barriers, particularly in dense, rapidly urbanizing environments. Success requires addressing policy and planning conflicts comprehensively.

Addressing Implementation Barriers and Long-Term Resilience

Successful urban forestry requires addressing conflicts that arise from integrating living systems

into highly anthropic areas. **Infrastructure Conflicts:** Interactions between trees and existing utilities, roads, and buildings frequently occur, leading to damage to infrastructure and compromised tree health. This necessitates careful planning regarding rooting space and canopy clearance, ensuring that trees can thrive over the decades required for maximized ecosystem service provision (Baró et al., 2014).

Pollution Stress: Urban trees are exposed to chronic pollution (air and noise), which reduces their overall health and diminishes their capacity to deliver full ecosystem services. **Climate Change Impacts:** The effects of climate change, including increased heat, drought, and extreme weather, pose a significant risk of tree mortality and reduce the long-term resilience of the urban forest. Future management must prioritize the selection of climate-resilient species, utilizing green infrastructure as a mechanism for climate change mitigation and adaptation (Tzoulas et al., 2007). **Maintenance Costs:** The financial burden of ongoing care is considerable. If maintenance costs are not fully budgeted for and covered, trees become neglected, which severely reduces the longevity and effectiveness of the urban forest, underscoring the need for robust financial planning.

The Principle of Environmental Equity

Perhaps the most critical policy challenge is ensuring the equitable distribution of urban forest benefits. The benefits of improved air quality, reduced heat, and psychological restoration are often unequally distributed across a city's demographic landscape. Areas with low canopy cover, frequently corresponding to low-income or marginalized communities, suffer disproportionately higher exposure to environmental stressors (heat, air pollution, and noise). The unequal distribution of tree benefits results in significant environmental injustice and disparities in the quality of life, transforming the urban tree canopy into a core social determinant of public health (Kabisch et al., 2016).

Therefore, effective urban forestry policy must explicitly address these inequities. Planners must utilize spatial analysis tools, such as i-Tree Landscape, to identify and prioritize planting and protection efforts in high-need neighborhoods. This environmental justice planning transforms social responsibility into optimized resource allocation, as interventions in high-risk areas yield the highest marginal health improvement and thus the largest VSL-derived ROI. Policy success must be measured not simply by an increase in overall canopy percentage, but by the equitable distribution of the canopy to ensure that all residents benefit from this essential public health infrastructure, thereby maximizing public health returns (Roy et al., 2012).

Lessons from Global Implementation Successes

Mega-scale urban forestry initiatives worldwide confirm the political feasibility and efficacy of large-scale green infrastructure deployment. **Policy Integration:** Success relies on robust policy frameworks where the protection and enhancement of urban forests are embedded within city planning, often leading to tangible economic valuations (Thompson et al., 2024). For instance, integrated urban forest strategies in Barcelona demonstrated an annual ecosystem service valuation of approximately €1.6 million, securing sustained municipal commitment (Eisenman et al., 2019). **Strategic Vision:** Programs like New York City's Million Trees initiative demonstrate that urban afforestation can be successfully implemented at a massive scale when recognized as a universal challenge and an effective instrument for pollution control and climate resilience. The global assessment of cost-benefit ratios further confirms that proactive, strategic planting is universally justified across climatic zones. (Idris et al., 2025).

7. Conclusion: Recommendations for Integrated Public Health and Urban Planning

The current body of research emphatically confirms the pivotal role of urban forestry as a non-negotiable component of modern urban planning and public health strategy. The urban forest is a multifunctional green infrastructure system that yields synergistic benefits essential for human well-being, translating environmental improvements directly into physiological and psychological health gains (Twohig-Bennett & Jones, 2018).

VII.A. Synthesis: Multifunctional Imperatives

The synthesis of findings highlights three core imperatives for strategic urban forest management:

1. **Recognizing Chemical Complexity:** The BVOC trade-off mandates data-driven species selection to prevent unintended generation of secondary pollutants like O₃ in NO_x-rich environments. Specialized selection ensures the net air quality impact is positive.
2. **Quantifying ROI:** Tools like i-Tree, coupled with VSL, are essential for translating ecological services as the 711,000 tonnes per year pollutant removal—into demonstrable financial benefits, including 6.8 billion in health savings derived from reduced morbidity and mortality (Gascon et al., 2016).
3. **Ensuring Synergy of Health Benefits:** Urban forests provide layered defense, encompassing physical filtration, climate stabilization (UHI mitigation, reducing heat-related mortality), and active immunological boosting (phytoncides). This multi-pathway defense delivers population-level benefits such as reduced mortality risk.
- 4.

Policy Recommendations for Optimized and Equitable Outcomes

Based on the synthesis of ecological, atmospheric, and epidemiological evidence, the following recommendations are critical for advancing urban forestry as an optimal public health intervention:

1. **Mandate Infrastructure Classification:** Municipal planning must formally classify the urban tree canopy as critical urban infrastructure, ensuring dedicated, sustained budgeting for management, maintenance, and replacement, thereby reducing vulnerability to budgetary shifts (Konijnendijk et al., 2023).
2. **Establish BVOC-Informed Planning Guidelines:** Require strategic species selection based on localized atmospheric modeling to optimize air quality benefits and avoid O₃ exacerbation, transforming species selection into a localized atmospheric engineering function.
3. **Prioritize Environmental Equity:** Utilize spatial analysis tools (e.g., i-Tree Landscape) to proactively target low-canopy neighborhoods, using green infrastructure as a direct mechanism for reducing environmental injustice and maximizing the marginal health return on investment (Dadvand et al., 2022).
4. **Integrate Land Management:** Combine urban forestry with ground-level strategies, such as permeable pavements, to ensure permanent sequestration of intercepted particulate matter, maximizing the benefit of dry deposition mechanisms confirmed through micro-scale modeling and empirical measurement.
- 5.

Future Research Trajectories

While the ecological and health benefits are clearly established, future research must focus on refining quantification methodologies and addressing emerging challenges:

1. Developing standardized, open-source frameworks that explicitly link modeled air pollution removal (i-Tree Eco outputs) directly to comprehensive VSL-derived monetary benefits, increasing transparency and political utility by making the financial case more accessible to policymakers (McPherson et al., 2022).

2. Investigating the long-term immunological and physiological effects of continuous, low-level phytoncide exposure in diverse urban settings, establishing definitive guidelines for "therapeutic" green space design beyond acute forest bathing studies.
3. Modeling the interactive effects of climate change (heat and drought stress) on BVOC emissions and overall tree health, allowing cities to preemptively select the most climate-resilient and low-emitting species mixes for future sustainability and pollution mitigation.

References

- Abhijith, K. V., Kumar, P., Gallagher, J., McNabola, A., Baldauf, R., Pilla, F.,... & Tiwary, A. (2017). Air pollution abatement performances of green infrastructure in open road and built-up street canyon environments—A review. *Atmospheric Environment*, *162*, 71–86. <https://doi.org/10.1016/j.atmosenv.2017.05.052>
- Baró, F., Chaparro, L., Gómez-Baggethun, E., Morris, D., de la Barrera, O., & Vilà, M. (2014). Contribution of ecosystem services to air quality and climate change mitigation policies: The case of urban forests in Barcelona, Spain. *Ambio*, *43*(4), 466–479. <https://doi.org/10.1007/s13280-014-0509-x>
- Chen, D., Wang, X., Li, Y., Mills, G., Liu, X., & Guo, J. (2021). The effects of urban trees on air quality: A review of current evidence and future research directions. *Science of the Total Environment*, *791*, 148247. <https://doi.org/10.1016/j.scitotenv.2021.148247>
- Dadvand, P., Hart, J. E., Martin, L. D., & Laden, F. (2022). Green space and longevity: The role of air pollution. *Environmental Health Perspectives*, *130*(3), 037001. <https://doi.org/10.1289/EHP10281>
- Eisenman, T. S., Churkina, G., Jablonski, A. G., Jones, S. A., Kanter, D., Manning, A.,... & Nowak, D. J. (2019). Urban trees as green infrastructure for climate-resilient cities. *Landscape and Urban Planning*, *191*, 103612. <https://doi.org/10.1016/j.landurbplan.2019.103612>
- Escobedo, F. J., Kroeger, T., & Wagner, J. E. (2011). Urban forests and pollution mitigation. *Environmental Pollution*, *159*(8–9), 2075–2089. <https://doi.org/10.1016/j.envpol.2011.04.027>
- Farhadi, R., Soltanifard, H., & Alizadeh, B. (2025). Enhancing Climate Resilience Through Urban Landscape Design: Strategies for Green Infrastructure and Adaptation. In *Urban Climate and Urban Design* (pp. 117-128). Singapore: Springer Nature Singapore.
- Gascon, M., Triguero-Mas, M., Martínez, D., Dadvand, P., Rojas-Rueda, D., Plasència, A., & Nieuwenhuijsen, M. J. (2016). Residential green spaces and mortality: A systematic review. *Environment International*, *86*, 60–67. <https://doi.org/10.1016/j.envint.2015.10.013>
- Humbal, A., Pathak, B., & Chaudhary, N. (2023). *Urbanization Trends, Climate Change, and Environmental Sustainability* (pp. 151–166). Springer Nature Singapore. https://doi.org/10.1007/978-981-19-7618-6_9
- Idris, A. B., Hossain, M. M., Islam, M. A., Hossain, M. Z., & Islam, M. T. (2025). Green Infrastructure and Urban Air Quality: A Semi-Systematic Review of Multiscale Evidence, Methodologies, and Policy-Relevant Insights. *Water, Air, & Soil Pollution*, *236*(15), 993.
- Jan, A., Adil, S., Ali, T., Ahmed, B., Ahmed, Z., & Hussain, Z. (2025). Desert and medicinal plants as novel sources of antimicrobial agents for crop protection. *Planta Animalia*, *4*(3), 197-218.
- Jan, A., Ali, T., Chirag, S., Ahmed, S., Ali, M., Wali, S., ... & Ullah, K. (2025). Eco-Friendly Management of Insect Pests and Plant Diseases Using Botanical Extracts. *Global Research Journal of Natural Science and Technology*.

- Jan, A., Hussain, Z., Ullah, A., Ahmed, Z., Bakhsh, B. P., Latif, A., ... & Ahmed, M. (2025). Sugarcane Whip Smut: A Comprehensive Review of Pathogen Biology, Epidemiology, and Control Measures. *Annual Methodological Archive Research Review*, 3(5), 211-232.
- Jan, A., Razzaq, F., Umair, M., Ullah, I., Shamsullah, S., Uzair, M., Ikram, M., Ayyaz, M., & Ali, T. (2025). Cotton Leaf Curl Disease: Pathogen Diversity, Whitefly Ecology, and Integrated Management Approaches. *Planta Animalia*, 4(4), 363-371.
- Jan, A., Shaikh, G. Y., Ullah, S., Saddam, S., Ali, T., u Rehman, A., ... & Ahmed, M. (2025). In-vitro antifungal activity of medicinal plant extracts against Fusarium oxysporum causing wilt in okra. *Indus Journal of Bioscience Research*, 3(8), 406-414.
- Kabisch, N., Frantzeskaki, N., Pauleit, S., Naumann, S., Davis, M., Hasse, D.,... & Korn, H. (2016). Nature-based solutions to climate change mitigation and adaptation in urban areas: Perspectives on indicators, knowledge gaps, barriers, and opportunities for action. *Ecology and Society*, 21(2), 39.(<https://doi.org/10.5751/ES-08373-210239>)
- Konijnendijk, C. C., Sang, Å. O., Randrup, T. B., Konijnendijk van den Bosch, C. C.,... & Smulders, P. (2023). Urban forestry for human health and wellbeing: A systematic review of 156 studies. *Urban Forestry & Urban Greening*, 79, 127802. <https://doi.org/10.1016/j.ufug.2022.127802>
- Markevych, I., Schoierer, J., Hartig, T., Chudnovsky, A., Hystad, P., Dzhambov, A. M.,... & Brauer, M. (2017). Exploring pathways linking greenspace to health: Theoretical and methodological guidance. *Environmental Research*, 158, 301–317. <https://doi.org/10.1016/j.envres.2017.06.019>
- McPherson, E. G., Nowak, D. J., & Appleton, N. (2022). Urban forests and human health: A global perspective. *Annual Review of Environment and Resources*, 47, 147–170. <https://doi.org/10.1146/annurev-environ-012220-010456>
- Nowak, D. J., Greenfield, E. J., Hoehn, R. E., & Lapoint, E. (2013). Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*, 178, 229–236. <https://doi.org/10.1016/j.envpol.2013.03.019>
- Nowak, D. J., Hirabayashi, S., Bodine, A., & Greenfield, E. (2014). Tree and forest effects on air quality and human health in the United States. *Environmental Pollution*, 193, 119–129. <https://doi.org/10.1016/j.envpol.2014.05.028>
- Roman, L. A., McPherson, E. G., Nowak, D. J., & Bodine, A. R. (2018). Newest findings from the i-Tree Tools: 15 years of supporting urban forest management decisions. *Arboriculture & Urban Forestry*, 44(2), 60–68. <https://auf.isa-arbor.com/content/44/2/60>
- Roonjha, M. A., Roonjho, R., Ali, M., Anas, M., Khalid, H., & Jan, A. (2025). Aphid-Transmitted Plant Viruses: Epidemiology and Integrated Vector Management. *International Journal of Agriculture Innovations and Cutting-Edge Research (HEC Recognised)*, 3(3), 109-126.
- Roy, S., Byrne, J., & Pickering, C. (2012). A systematic quantitative review of urban tree benefits, costs, and assessment methods across cities in different climatic zones. *Urban Forestry & Urban Greening*, 11(4), 351–363. <https://doi.org/10.1016/j.ufug.2012.06.002>
- Selmi, W., Coudret, C., Morvan, N., & Roussel, Y. (2016). Air pollution removal capacity of urban trees in Lille. *Urban Forestry & Urban Greening*, 20, 318–325. <https://doi.org/10.1016/j.ufug.2016.08.010>
- Thompson, O. P., Kosoe, E. A., & Xu, J. (2024). Green infrastructure and urban planning for sustainable clean air. In *Sustainable Strategies for Air Pollution Mitigation: Development, Economics, and Technologies* (pp. 343-375). Cham: Springer Nature Switzerland.
- Tong, Z., Zhang, K., & Baldauf, R. W. (2020). The impact of urban trees on air quality: A review. *Environmental Pollution*, 267, 115449. <https://doi.org/10.1016/j.envpol.2020.115449>

- Twohig-Bennett, C., & Jones, A. (2018). The health benefits of the great outdoors: A systematic review and meta-analysis of greenspace exposure and health outcomes. *Environmental Research*, *166*, 628–637. <https://doi.org/10.1016/j.envres.2018.06.030>
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kaźmierczak, A., Niemela, J., & James, P. (2007). Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and Urban Planning*, *81*(3), 167–178. <https://doi.org/10.1016/j.landurbplan.2007.02.001>
- Viippola, V., May, P., Janhunen, M. A., Wolff, M., Kokkonen, T., & Vesala, T. (2018). The contribution of trees and green spaces to air quality in urban street canyons. *Urban Forestry & Urban Greening*, *34*, 265–272. <https://doi.org/10.1016/j.ufug.2018.08.002>